

## PATENT COOPERATION TREATY

PCT

## NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

Assistant Commissioner for Patents  
United States Patent and Trademark  
Office  
Box PCT  
Washington, D.C. 20231  
ÉTATS-UNIS D'AMÉRIQUE

in its capacity as elected Office

Date of mailing (day/month/year) 30 January 2000 (30.01.00)	
International application No. PCT/SE99/00824	Applicant's or agent's file reference 2998459
International filing date (day/month/year) 12 May 1999 (12.05.99)	Priority date (day/month/year) 14 May 1998 (14.05.98)
Applicant LINDAHL, Åke et al	

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International Preliminary Examining Authority on:

09 December 1999 (09.12.99)

☐ in a notice effecting later election filed with the International Bureau on:2. The election ☒ was☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland	Authorized officer Martine Lee
Facsimile No.: (41-22) 740.14.35	Telephone No.: (41-22) 338.83.38

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>A61K 9/00, 9/70</b>		(11) International Publication Number: <b>WO 99/58109</b>	
A1		(43) International Publication Date: 18 November 1999 (18.11.99)	
(21) International Application Number: PCT/SE99/00824		<b>(81) Designated States:</b> AE, AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 12 May 1999 (12.05.99)			
(30) Priority Data: 9801705-6 14 May 1998 (14.05.98) SE			
(71) Applicant (for all designated States except US): BIOGLAN AB [SE/SE]; P.O. Box 50310, S-202 13 Malmö (SE).			
(72) Inventors; and (75) Inventors/Applicants (for US only): LINDAHL, Åke [SE/SE]; Ringduvevägen 50, S-274 33 Skurup (SE). HAGSLÄTT, Håkan [SE/SE]; Flygelvägen 14, S-224 72 Lund (SE). HEIMAN, Catharina [SE/SE]; Kilian Zollsgatan 11A, S-217 56 Malmö (SE). BRYLAND, Rickard [SE/SE]; V. Rönneholmsvägen 60B, S-217 41 Malmö (SE).			
(74) Agent: AWAPATENT AB; P.O. Box 45086, S-104 30 Stockholm (SE).		<b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: BIOLOGICALLY ACTIVE COMPOSITION			
(57) Abstract <p>The invention relates to a novel biologically active composition which comprises a biologically active agent to be released therefrom, said biologically active agent being dissolved and/or dispersed in a supersaturated state within a carrier, which carrier is a liquid and/or solid non-crystalline matrix, and where the precipitation of said biologically active agent is substantially, or completely, inhibited therein. Said supersaturated state is obtainable by subjecting one or more carrier starting substance(s) to such chemical operation(s) that a matrix is provided in which the degree of saturation of said biologically active agent is higher than the degree of saturation of said biologically active agent in said carrier starting substance(s), the biologically active agent being added either before said chemical operation(s) or after a predetermined point of time, after which the composition thus prepared is further subjected to said chemical operation(s).</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GHI	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

BIOLOGICALLY ACTIVE COMPOSITIONTechnical Field

The present invention relates to a biologically active composition from which one or more biologically active components are to be released. More specifically, the invention relates to a biologically active composition wherein the biologically active agent is present in a supersaturated state within a carrier without being precipitated therefrom.

10 Background of the Invention

From *inter alia* toxicological points of view, it is often preferred, upon treatment of diseases or symptoms thereof, to deliver drugs directly to their site(s) of action. It is well known that the risks of obtaining detrimental effects of systemic origin are often markedly reduced if a drug is delivered directly to its site(s) of action. Furthermore, systemic delivery often involves metabolism of the drug prior to its appearance at the site of action, which leads to a subsequent reduction of its biological effect. Another important aspect is that in e.g. cases of imminent overdose, allergic reactions or administration of contraindicating drugs, it is easy to remove topical compositions in contrast to drugs administered per-orally or by injection.

25 As used herein, topical administration comprises *inter alia* dermal, sub-lingual, gingival, buccal, transdermal, nasal, vaginal and rectal administration, whereby the resulting biological effect may be local and/or systemic.

30 In e.g. dermal, nasal, vaginal, buccal or sub-lingual administration, only a very limited number of drugs are capable of permeating into the human body by themselves at a useful rate. Consequently, a lot of research has been conducted in order to investigate the possibility of both improving traditional non-invasive

35

delivery techniques and developing novel non-invasive drug delivery systems or devices intended for systemic and/or internal use. Three fundamentally different approaches towards this objective have been disclosed.

5        Firstly, there is the well known possibility of improving the penetration properties of the drug by chemical modification thereof. After the drug has entered the body, its pharmacologically active form is obtained by chemical reaction(s) *in vivo*. However, this so called  
10        pro-drug approach is only occasionally a succesful alternative. There are several reasons therefor, such as i) the penetration rate of the pro-drug may still be too low, ii) the pro-drug may be toxic or otherwise harmful, or iii) the *in vivo* conversion to the active form of the  
15        drug is too slow and/or partially results in inactive or toxic compounds. A distantly related approach is the preparation of an ion pair between a drug and an appropriate counter ion. However, generally such an ion pair does not display any markedly improved penetration  
20        rate through human barriers.

      Secondly, the properties of the barrier may be changed in order to facilitate the drug delivery. Methods of achieving this are e.g. ultra-sonication, applying of electrical current or the use of so called penetration  
25        enhancers in the composition. All of these methods act by disrupting the structure of the barrier, thereby facilitating drug diffusion through the barrier into the body, and/or improving the drug solubility in the barrier. However, the methods involving e.g. heat, ultra-  
30        sonication and electrical current are generally not designed for being easily managed by the patient in a convenient manner, and therefore require hospitalisation, which is a major disadvantage with said methods. In  
35        addition, all methods which are based on the approach of changing the barrier properties are questionable from a toxicological point of view due to the observations that i) adverse effects on the cells of the barrier have been

demonstrated, and ii) a reduction of the protective properties of the barrier also result in increased penetration rate for any substance, not only the drug, that is present at the site of administration. It should also be mentioned, that a majority of the known chemical penetration enhancers require some time for the onset of their action, i.e. display a lag time of action, since they must be established in the barrier before the actual increase in penetration rate is observed.

10 Thirdly, the driving force of the drug for entering the body can be changed. That is, the difference in the electrochemical potential of the drug between the drug reservoir and the body can be increased. Drug delivery systems based on this approach result in a high flux of the drug through the barrier and usually also display a reduced lag time of action.

In methods based on iontophoresis, this approach is utilised by applying an electrical potential gradient across the barrier. Obviously, these methods are mainly suitable for drugs having a net charge and are therefore much less efficient for uncharged and zwitterionic species, since the flux of the two latter species is improved mainly due to e.g. osmotic and electroosmotic driving forces. Iontophoresis methods also have the disadvantage that they may alter the structure of the barrier.

In another approach, the flux of a drug into the body can be enhanced by increasing the chemical potential of the drug in the carrier therefor. This is normally performed by chemical optimisation of the drug composition by adjusting the degree of saturation of the drug in said carrier. The methods based on this approach offer several advantages as compared to the previously mentioned methods, since the flux of the drug is increased in comparison with subsaturated and saturated systems. Furthermore, the properties of the barrier itself are comparatively less affected and the lag time

of initiation for the pharmacological effect is reduced. There are two particularly important aspects in this approach:

- i) creation of an initial high chemical potential of the drug in the composition
- ii) maintenance of a high chemical potential of the drug in the vicinity of the barrier after the application of the composition.

Therefore, it is usually desirable to prepare pharmaceutical compositions which are saturated with respect of the drug. During application, another important aspect of said composition is that the solubility and diffusion properties of the drug in the used vehicle must preclude depletion of the drug in the vicinity of the barrier. Examples of compositions used for this purpose are microemulsions and emulsions.

Another approach towards keeping the composition saturated is the use of an excess amount of drug (non-solubilised) in the carrier, whereby the drug is subsequently dissolved as it replaces the drug which has penetrated through the barrier.

Yet another approach is the use of a supersaturated composition of the drug. Here, the driving force of the drug to penetrate the barrier is higher than in the saturated composition, since the drug in a supersaturated composition has higher chemical potential in comparison with the corresponding saturated composition. For example, such compositions have been prepared according to the following means or principles: i) dissolving the drug at temperatures and/or pressures at which the solubility of the drug is higher as compared to those temperatures and/or pressures that are relevant for medication (W.L. Chou and S. Riegelmann, *J. Pharm. Sci.*, Vol.60, No.9, pp.1281-1302, 1971; WO 97/10812), ii) using solid dispersions or eutectic mixtures or solid drug particles of low degree of crystallinity or of high energy polymorphs (W.L. Chou and S. Riegelmann, *supra*),

iii) mixing a saturated drug solution with a non-solvent therefor, thereby performing a merely physical operation, in situ or prior to application, with or without the presence of an antinucleating agent (US 4 940 701; 5 US 4 767 751), iv) solvent evaporation to the surrounding air (Coldman et al., *J. Pharm. Sci.*, 58, No.9 (1969), pp 1098-1102), v) solvent penetration into the human body, vi) water uptake into the composition from the human body, vii) pH-changes in the composition caused by H<sup>+</sup>- 10 uptake from the human body, or viii) dispersing an aqueous solution or emulsion of a drug in an aqueous dispersion of a polymer latex (Lichtenberger et al., *"Polymer films from aqueous polymer dispersions as carriers for transdermal delivery of lipophilic drugs"*, 15 15th Int Symp CRS:Basel 1988; Abstr 89). An important common denominator of iv)-vii) is that the supersaturation is not initially present in the composition, and is therefore de facto not accomplished until the composition is applied to a human body. 20 Furthermore, a major problem with all the compositions i)-viii) is that the drug generally precipitates in a relatively short time, in which case the saturation degree becomes markedly reduced.

In DD 217 989, a subsaturated solution of a drug is 25 mixed with a solution or suspension of an acrylate, after with the mixture so prepared is dried, whereby a supersaturated composition is obtained by use of an exclusively physical operation.

W.L. Chou and S. Riegelmann (*J. Pharm. Sci.*, Vol.58, 30 No. 12, pp.1505-1510, 1969) have reported that in matrices of higher molecular weight polyethylene glycols, precipitation of a supersaturated drug dissolved therein is usually sluggish. In said document, supersaturation was obtained through either direct melting or solvent 35 concentration, i.e. by use of typical physical operations.



As prior art, reference is also made to WO 97/00670, which discloses a composition based on ingredients similar to those utilized in the present invention. However, said reference does not disclose or suggest any  
5 supersaturated state or even less those features and measures of the present invention which have been found crucial to impart a stable, supersaturated state to such a composition.

Other prior art of interest is WO 97/10812, which  
10 discloses a method for preparing supersaturated systems, wherein an admixture of drug and polymer having a calculated depressed melting temperature is heated to a temperature above said calculated temperature, whereby the drug is dissolved in the polymeric material and  
15 supersaturation thereof is obtained through cooling of the heated solution. However, the present invention is not related to preparation of supersaturated systems by exploitation of the calculated depressed melting temperature of an admixture through an entirely physical  
20 operation.

Mention can also be made of GB 2 306 885, which utilises the skin's innate ability to buffer applied liquids. Here, a supersaturated system is attained in situ by applying a subsaturated drug composition having a  
25 pH of 7-12 or 3-4 to skin, where the buffering effect of skin causes a pH change to 4.5-6.5, whereby a supersaturated composition is obtained by means of a change of the degree of protonation of the drug. The preparation of supersaturated systems according to the  
30 present invention does not rely on such an exchange of protons.

#### General Disclosure of the Invention

A novel approach for obtaining a biologically active composition with outstanding delivery rate of its active  
35 component(s) has now been developed, wherein said composition comprises a biologically active agent which is present in a substantially stable supersaturated

state. In brief summary, it has been found that by  
subjecting a carrier starting substance to such chemical  
operation(s) that a carrier matrix of substantially non-  
crystalline or amorphous nature is created, in which the  
5 degree of saturation of a biologically active agent is  
higher than the degree of saturation of said agent in the  
starting carrier substance, a surprisingly stable  
supersaturated composition can be obtained. In the  
composition thus prepared, the precipitation of said  
10 agent is substantially, or completely, inhibited by said  
carrier matrix *per se*.

The term "biologically active agent", as used  
herein, also comprises such progenitors thereto which are  
readily transformable, e.g. enzymatically and/or  
15 hydrolytically, to a biologically active agent *per se*.

Thus, the present invention relates to a novel  
biologically active composition which comprises a  
biologically active agent to be released therefrom, said  
biologically active agent being dissolved and/or  
20 dispersed in a supersaturated state within a carrier,  
which carrier is a liquid and/or solid substantially non-  
crystalline matrix, and where the precipitation of said  
biologically active agent is substantially, or  
completely, inhibited therein.

25 The term "liquid" as used in connection with the  
present invention should be interpreted in a broad sense,  
viz as any material being a mobile or viscous liquid,  
rubber, glass or plastic; thus including solutions,  
creams, pastes, ointments and gels within the scope of  
30 the claims.

The present invention also relates to a method for  
the preparation of a biologically active composition  
comprising a biologically active agent dissolved and/or  
dispersed in a supersaturated state in a carrier therefor  
35 as well as to said composition for use as a medicament.

The term "pharmaceutically active agent", as used  
herein, also comprises such progenitors, e.g. pro-drugs,

which are readily transformable, e.g. enzymatically and/or hydrolytically, to a pharmaceutically active agent *per se*.

One of the objects of the present invention is thus  
5 to provide a supersaturated composition which does not display any significant precipitation or loss of effect during long-term storage at room temperature, or even at above or below room temperature, during e.g. months or even years.

10 Another object of the present invention is to provide a supersaturated composition which does not display any significant precipitation or loss of effect during its application to a human or animal patient.

Still another object of the present invention is to  
15 provide a carrier matrix which is suitable in preparation of a composition having a particularly high degree of supersaturation of a drug (*vide infra*).

Yet another object is to provide a stable  
supersaturated composition which is easily handled and  
20 does not require professional assistance upon use thereof.

As a result of the high delivery rate of its active component(s), another object of the present invention is to provide a composition which allows for efficient  
25 topical treatment, preferably dermal or transdermal administration to small areas, which is a general advantage in the topical administration of drugs.

#### Detailed Disclosure of the Invention

30 More specifically, the invention refers to a biologically active composition comprising a biologically active agent dissolved and/or dispersed in a carrier therefor, wherein said carrier is a liquid and/or solid substantially non-crystalline matrix in which said  
35 biologically active agent is present in a supersaturated state and in which the precipitation of said biologically active agent is substantially, or completely, inhibited

by said matrix, said supersaturated state being obtainable, or obtained, by subjecting one or more starting substance(s) to such chemical operation(s) that a liquid and/or solid substantially non-crystalline matrix is provided in which the degree of saturation of said biologically active agent is increased in comparison with the degree of saturation of said agent in the starting substance(s), the biologically active agent being added before said chemical operation(s) has (have) been completed.

As used herein, the term "chemical operation" refers to a measure resulting in formation or cleavage of covalent bonds. Said formation or cleavage may comprise or by indirect means yield a pH change of the composition, thus involving a proton transfer which in some cases may be regarded as formation or cleavage of a covalent bond. However, such a pH change is in this context the result of a chemical operation which does not merely comprise a proton transfer but which also comprises formation or cleavage of other types of covalent bonds.

In one embodiment of the invention, said supersaturated state is obtainable by subjecting one or more carrier starting substance(s) to such chemical operation(s) that a matrix is provided in which the degree of saturation of said biologically active agent is higher than the degree of saturation of said biologically active agent in said carrier starting substance(s), the biologically active agent being added at a predetermined point of time after said chemical operation(s) have been initiated, after which the composition thus prepared is further subjected to said chemical operation(s).

Other preferable embodiments of the composition claimed will be defined in the claims or referred to below in connection with the method.

Thus, the present invention also refers to a method for the preparation of a biologically active composition

comprising a biologically active agent dissolved and/or dispersed in a carrier therefor, wherein

a carrier starting substance, or a mixture of two or more different starting substances, is (are) subjected to such chemical operation(s) that a liquid and/or solid non-crystalline carrier matrix is formed, in which the degree of saturation of a biologically active agent is higher than the degree of saturation of said agent in said carrier starting substance(s), said biologically active agent being added before said chemical operation(s) has (have) been completed and in an amount such that a supersaturated state is obtained. Generally this means that said chemical operation(s) is (are) initiated either:

- i) in the presence of said biologically active agent; or
- ii) in the absence of said biologically active agent, after which said agent at a predetermined point of time is added and the composition thus prepared is further subjected to said chemical operation(s);

addition of said biologically active agent in both i) and ii) being made using an amount such that a supersaturated state is obtained.

In one embodiment of the invention, the degree of saturation of a biologically active agent is higher as a result of such chemical operation(s) that a liquid and/or solid non-crystalline carrier matrix is formed, in which the solubility of a biologically active agent is lower than the solubility of said agent in said carrier starting substance(s).

In another embodiment of the invention, the degree of saturation of a biologically active agent is higher as a result of such chemical operation(s) that a liquid and/or solid non-crystalline carrier matrix is formed, in which the degree of dissociation, aggregation and/or degree of protonation of a biologically active agent is

different from the degree of dissociation, aggregation and/or degree of protonation of said agent in said carrier starting substance(s). As a non-limiting example, this embodiment allows formation *in situ* of a suitably charged, e.g. protonated or deprotonated, or non-charged form of said biologically active agent, which form has a higher skin penetration rate in comparison with the form of said agent present before said chemical operation(s) is initiated.

10 In yet another embodiment of the invention, the degree of saturation of a biologically active agent is increased by such chemical operation(s) that both the two embodiments set forth above are practised either simultaneously or consecutively.

15 In one embodiment of the invention, said biologically active agent is being added, either above or around room temperature, in solid and/or liquid, i.e. melted, state and is subsequently dissolved in said starting substance(s) either above or around room temperature.

20 In another embodiment of the invention, said biologically active agent is being added, either above or around room temperature, as a solution or dispersion and is subsequently dissolved in said starting substance(s) either above or around room temperature.

25 According to the present invention, above room temperature is a temperature above about 25°C, such as about 25-200°C, preferably about 30-150°C. Examples of other suitable temperatures are about 35-100°C and 40-80°C.

30 The particular addition method used for said agent can be any common inclusion technique available to a person skilled in the art, and said solution or dispersion of the biologically active agent can be prepared *inter alia* by solvent evaporation, freeze-drying or by use of any one of the methods i)-vii) (*vide supra*).

Preferably, in the composition according to the invention as well as in the method for preparation thereof, the starting substance(s) act(s) as solvent or dispersing medium.

5 Said chemical operation(s) generally involve(s) one or more chemical reactions, preferably etherifying, esterifying, hydrolysis, substitution, addition, elimination, oligomerising and/or polymerising reactions, wherein polymerising reactions are the most preferred.

10 Said carrier starting substance(s), which is subsequently subjected to said operation(s) above, is selected from monomers, acids, such as mono-, di- or triacids or higher acids, alcohols, including mono-, di- or triols, ketones, aldehydes, amines, amides,  
15 anhydrides, lactides, glycolides, saccharides and derivatives thereof, acrylic or acrylamide type compounds, such as methyl methacrylate, monomers of PEO-diacrylate (PEO=polyethylene oxide), cyanoacrylate, acrylate saccharides, including acrylate starch, acrylate  
20 lactate, acrylate glycolate, isocyanates, ethylene oxide, propylene oxide, pyrrolidone, PEO-diacrylate, ethylene-vinyl acetate, monomers of organic siloxanes and oligomers, polymers or prepolymers thereof. As indicated earlier, one, two or more of the above substances can be  
25 chosen, thereby allowing the formation of co-polymers and/or higher polymers.

It is to be understood by a person skilled in the art, that said chemical operation(s) is performed to such a degree of completion that a desired non-crystalline  
30 carrier matrix is obtained, which matrix is optimal for a particular biologically active agent in a particular context. Thus, all of the starting substance(s) present when said chemical operation(s) is initiated do not necessarily have to react completely in order to carry  
35 out the invention, as long as the desired degree of supersaturation is attained.

In a preferred embodiment of the present invention, the carrier starting substances are an acid and an alcohol, said formed non-crystalline matrix comprising, or being, an ester and/or polyester thereof. In a more preferred embodiment, said carrier starting substances  
5 are citric acid and propylene glycol.

In an alternative embodiment, the starting substance is one bi- or multi-functional substance only, which when subjected to said chemical operation(s) provides the  
10 desired non-crystalline carrier matrix by chemical reaction(s) with itself. In a non-limiting disclosure, such a starting substance can be citric acid, which when subjected to esterifying conditions provides a non-crystalline citric acid ester and/or polyester matrix  
15 according to the invention.

According to the present invention, suitable chemical operation(s) involve(s) subjecting said carrier starting substance(s) to such polymerising conditions which are normally used, according to standard reference  
20 literature, for the selected starting substance(s) or combinations thereof. Furthermore, such polymerising conditions should be chosen in order to optimise the manufacturing procedure, in respect of e.g. the stability of said agent, manufacturing time and degree of  
25 supersaturation, for the particular biologically active agent used. Typically, said conditions comprise e.g. subjecting said carrier starting substance(s) to a temperature from around -50°C to around 300°C, preferably around 0-150°C. Other examples of useful temperature  
30 ranges are 20-100°C and 50-80°C. Said temperature ranges are particularly preferred when the starting substance(s) are a mixture of citric acid and propylene glycol. Naturally, said chemical reaction(s) are selected and performed so that in each case the maximum or optimum  
35 delivery rate of said biologically active agent is obtained.



Preferably, said chemical reaction(s) is (are) performed for a time period of from 1 minute to 6 months, more preferably from 0,5 hours to 4 months. As an example, said time period may also be from 1 hour to 3 months or from 1 to 2 months.

The predetermined point of time (*vide supra*), as measured after said chemical operation(s) has (have) been initiated, is generally from 1 minute to 6 months, preferably from 0,5 hours to 4 months, after which the composition thus obtained is further subjected to said chemical operation(s) for a time period of about from 1 minute to 6 months, preferably from 0,5 hours to 4 months. As an example, said predetermined point of time may also be from 1 hour to 3 months or from 1 to 2 months.

The used chemical reaction(s) in the present invention preferably comprise a polymerisation reaction and most preferably such reaction in which ether and/or ester bonds are formed. Other preferred polymerisation reactions are step polymerisation reactions and chain polymerisation reactions comprising either radical initiation, ionic initiation or coordination complex initiation.

According to the present invention, some of the monofunctional starting substance(s) above, e.g. monoacids and -alcohols, can also be used to form a non-crystalline matrix consisting of e.g. monoesters and monoethers. Monofunctional monomers can also be introduced into said chemical reaction as a means of modifying the reaction or controlling the end point thereof.

As already indicated, in order to efficiently inhibit precipitation of the supersaturated biologically active agent, said formed matrix is of a substantially non-crystalline, or amorphous, nature. Polymers, co-polymers, oligomers and ethers or esters of the previously

outlined starting substance(s) (*vide supra*) are particularly useful for this purpose.

A number of different parameters have been of interest in the development of the present invention. As an example of such a parameter, a reaction which results in the formation of a non-crystalline matrix, consisting of molecules with a larger molecular weight than the starting substance(s), can result in an increase of the thermodynamic potential of the form of the biologically active agent(s) which diffuse(s) through a biological barrier, such as skin. During the progression of such a reaction, a lowered solubility of the biologically active agent in said matrix will in many instances be observed, albeit it must here be emphasized that said lowered solubility may not always be necessary in order to yield an increased thermodynamic potential of the form of the biologically active agent which *de facto* diffuses through the skin. Moreover, the degree of dissociation, aggregation and/or protonation of the biologically active agent, e.g. as a result of pH changes, is often relevant in eliciting the desired increased thermodynamic potential of the form(s) of said agent diffusing through the skin.

Non-limiting examples of biologically active agents, preferably pharmaceutically active agents, which are suitable for use in the present invention are e.g., guanosides, corticosteroids, psychopharmaceutical hormones, oxicams, peptides, proteins as well as agents selected from the group of antibiotics, antivirals, antimicrobials, anticancer agents, antifungals, oestrogens, antiinflammatory agents, neuroleptic agents, melanocyte stimulants and gland stimulants, preferably stimulators of sebaceous and pilo-sebaceous glands, and agents with an effect on mast cell secretion.

In an alternative embodiment of the present invention, the biologically active agent may also react reversibly with said starting substance(s) in such a

manner that e.g. esters, ethers, co-polymers and/or other conjugates are formed. Thus, this embodiment allows preparation of a non-crystalline matrix containing both said biologically active agent in a substantially stable  
5 supersaturated state and conjugate(s) thereof, whereas said conjugate(s) can be present in either a subsaturated, saturated or supersaturated state. Alternatively, said conjugate(s) can be present in a supersaturated state, whereas said biologically active  
10 agent is present in either a subsaturated, saturated or supersaturated state. Therefore, in the case where said biologically active agent is a drug, this particular embodiment allows formation in situ of a corresponding drug progenitor, which may either function as a pro-drug  
15 or as a depot of the supersaturated drug, or a combination of both. As an example of this embodiment, a biologically active agent containing a carboxylic acid or alcoholic functionality may form an ester with said carrier starting substance(s) when a mixture thereof is  
20 subjected to esterifying conditions.

In another embodiment of the present invention, the starting substance(s) can be an ester and/or polyester matrix, or an ether and/or polyether matrix, to which a biologically active agent is added, after which the  
25 dispersion or solution formed is subjected to a hydrolysis reaction providing a liquid and/or solid non-crystalline carrier matrix in which the degree of saturation of said biologically active agent is higher than the degree of saturation of said biologically active  
30 agent in said starting substance(s), a stable supersaturated dispersion or solution thus being obtained. As a non-limiting example of such an embodiment, the starting substance(s) may consist of several esters and/or polyesters, of which one or several  
35 is much more readily hydrolysed in comparison with all other substances present, including the biologically active agent.

In yet another embodiment of the invention, a minor amount of said starting substance(s) is subjected to said chemical conditions, preferably a polymerisation, in the presence of a solvent, whereby a supersaturated one- or two-phase matrix is formed, such as a liquid/solid non-crystalline matrix.

However, in the most preferred embodiment, the biologically active composition consists of one liquid or solid phase only.

As earlier indicated, in another embodiment of the present invention the carrier starting substance(s) can be subjected to said chemical reaction(s), preferably a polymerisation, in advance and without the presence of said biologically active agent. By using this approach, a prefabricated liquid and/or solid non-crystalline matrix is provided, to which matrix a biologically active agent can subsequently be added at a predetermined point of time by use of any suitable inclusion method, such as e.g. mixing, heating, freeze-drying and/or solvent evaporation, after which the composition thus prepared is further subjected to said chemical reaction(s), which is (are) either identical or somewhat modified, by e.g. use of a lower reaction temperature or further addition of one or more of the previously outlined starting substance(s), in comparison with the chemical reaction(s) performed initially.

For some biologically active agents it is preferred to prepare a supersaturated composition shortly before administration thereof. Indeed, the present composition is useful for such preparations in addition to it being suitable for supersaturated compositions intended for long-term storage and application. As for the choice of a suitable degree of supersaturation of the biologically active agent in the present composition, it is known from the laws of thermodynamics that within a given period of time the danger of precipitation increases with the degree of supersaturation. Still, the present composition

is also suitable in such particular preparations where a very high degree of supersaturation is desirable, despite a somewhat increased danger of precipitation.

The scope of the present invention is not limited to  
5 the specific embodiments disclosed above, and the  
disclosed invention may optionally be combined with the  
methods i)-vii) (*vide supra*) in any suitable manner, if  
deemed necessary in any particular case. As a non-  
limiting example, the pH of the composition prepared  
10 according to the invention may optionally be subsequently  
modified by inclusion of a suitable acidic or basic  
compound, if useful in a particular context.

The following non-limiting example will illustrate  
the present invention further.

Brief description of the enclosed diagrams

Diagram 1 shows the amount of permeated metronidazole as a function of time for a subsaturated composition  $A_0$ , a saturated composition C and the  
5 supersaturated compositions  $B_1$  and  $B_2$ .

Diagram 2 displays the amount of metronidazole permeated from compositions X1-X4 and Y1-Y4.

Experimental part

10 Example 1; demonstrating an increased thermodynamic potential by using the method of the present invention:

The degree of supersaturation was characterised by the permeation rate of the biologically active agent through a membrane (Silastic sheeting NRV, 0,005 inches, serial #HH055353) by using a Franz diffusion cell (FDC-  
15 400 Crown Glass Company) with a cell opening area of 2,011 cm<sup>2</sup>. All permeation rate measurements were performed at 25°C and deaerated H<sub>2</sub>O was used as acceptor phase on the opposing side of the membrane. The donor and  
20 acceptor phase were both sealed with parafilm, and each experiment was performed in triplicate.

Starting substances: citric acid (CiAc) and propylene glycol

Four parts of CiAc and six parts of propylene glycol  
25 were added to a sealable container at room temperature, after which said container was sealed. The resulting mixture was stirred with a magnetic stirrer and the temperature was raised to and maintained at 80°C until all CiAc was dissolved, after which the solution was  
30 allowed to attain room temperature. This solution was denoted A. Solid metronidazole was then added to the solution A in a 5:95 ratio (w/w), after which the metronidazole was dissolved by magnetic stirring at room temperature. The solution thus prepared was then split  
35 into two solutions denoted  $A_0$  and B, respectively.

As reference, a solution of 4 parts of CiAc and 6 parts of propylene glycol was prepared as above. Solid metronidazole in a 7,5:92,5 ratio (w/w) was added, and the mixture was stirred at room temperature for three days. After centrifugation resulting in sedimentation of non-dissolved metronidazole, the obtained supernatant thus consisted of a saturated metronidazole composition, denoted C. The obtained final ratio between metronidazole and CiAc/propylene glycol was 7:93 (w/w).

The underlying principles behind the compositions A-C were the following:

$A_0$  is a subsaturated mixture of a pharmaceutically active agent and carrier starting substances which is not actively subjected to polymerisation;

in B, the starting substances are subjected to polymerisation conditions in the presence of a pharmaceutically active agent; and

in C, the permeation rate for a saturated solution of a pharmaceutically active agent in a matrix of said carrier starting substances is illustrated.

The compositions B and C were then treated as follows:

B was split into two compositions, which were stored at 70°C for one month ( $B_1$ ) and two months ( $B_2$ ), respectively, after which time period the permeation rate measurements were performed on the formed compositions  $B_1$  and  $B_2$ , respectively.

The compositions  $A_0$  and C were used directly after the preparation thereof.

The measured permeation rates are depicted in the enclosed Diagram 1.

Diagram 1 shows that a considerably higher permeation rate is obtained in the compositions  $B_1$  and  $B_2$ , as compared to any one of the compositions A or C. This increased permeation rate is in turn clear evidence that the thermodynamic potential of metronidazole is significantly higher in the compositions  $B_1$  and  $B_2$  in

comparison with any one of the compositions A<sub>0</sub> or C. Here, it is important to note that the compositions A<sub>0</sub> and B are initially the same.

In summary, this example shows that supersaturation  
5 of the initially subsaturated composition is attained upon polymerisation. Indeed, further polymerisation results in an even higher permeation rate, i.e. a higher thermodynamic potential, as is illustrated by B<sub>1</sub> and B<sub>2</sub>.

10 Example 2; demonstrating the precipitation preventing properties of the carrier matrix of the present invention:

A Franz diffusion cell as disclosed above was used under conditions similar to those of example 1, unless  
15 otherwise noted. Permeation rate experiments were performed for 21 h. As a reference, the permeation rate from the saturated composition C in example 1 was determined to be 46 µg per 21 h in a Franz diffusion cell experiment, as depicted in Diagram 1. The experiments  
20 were all analysed by use of spectrophotometry. The results are depicted in diagram 2.

In order to determine its water solubility, an excess of metronidazole was added to water, after which the mixture was stirred for 3 days at room temperature.  
25 Analysis by spectrophotometry was performed after sedimentation and centrifugation, and a resulting solubility of s=0.82% (w/w) was obtained.

Four supersaturated water solutions of metronidazole were then manufactured, each one having a degree of  
30 saturation (DS = concentration/solubility) of 1.3, 1.6, 2.0 and 2.5, respectively. They were prepared by heating the corresponding amount of metronidazole in water to 80°C for 30 min under stirring, followed by equilibration to room temperature, thereby yielding supersaturated  
35 solutions. The time for precipitation of metronidazole to occur (t<sub>p</sub>) upon storage at room temperature was monitored



by visual inspection, and the results are shown in Table 1.

Table 1. Time for precipitation of metronidazole from a supersaturated solution thereof in water.

Solution	metronidazole conc. % (w/w)	DS*	$t_p$
1	1.06	1.3	5 days < $t_p$ < 14 days
2	1.31	1.6	2 h < $t_p$ < 17 h
3	1.65	2.0	3 h < $t_p$ < 3.5 h
4	2.05	2.5	0.5 h < $t_p$ < 1 h

\*DS=1 equals 0.82 % (w/w) metronidazole in water (vide supra), as determined by spectrophotometry

10 A composition X was manufactured by mixing 4 parts of CiAc and 6 parts of propylene glycol (starting substances) at room temperature in a glass container which was subsequently sealed. The temperature was raised to and maintained at 80°C under stirring for about 45  
15 min. The resulting solution was kept at room temperature for about 30 min, and was then split into 4 separate solutions. An appropriate amount of metronidazole (see Table 2) was subsequently added to each solution, followed by heating of the mixture at 80°C for about 40  
20 min, after which the resulting compositions were allowed to attain room temperature, thereby yielding the supersaturated compositions X1-X4. Directly after their preparation, the compositions X1-X4 were investigated by Franz diffusion cell measurements (see example 3).

25 A sample was taken from each respective composition X1-X4. These four samples were each kept at 70°C for 3 weeks, thereby yielding the compositions Y1-Y4 (see Table 2). Y1-Y4 were also examined in Franz diffusion cell experiments (see example 3).

Table 2. Degree of saturation of metronidazole in the compositions X1-X4 and Y1-Y4.

metronidazole conc. % (w/w)	Composition	DS*	Composition	DS*
8.0	X1	1.16	Y1	1.62
9.0	X2	1.23	Y2	1.86
10.0	X3	1.54	Y3	2.02
11.0	X4	1.59	Y4	2.18

\*The permeation rate at saturation was assumed to be  
5 46 µg per 21 h.

The DS values shown in Table 2 were obtained by use  
of Franz diffusion cell measurements, and to a person  
skilled in the art, it is well known that the permeation  
10 rate of a compound through a Silastic membrane in a Franz  
cell diffusion cell experiment is a direct measure of the  
thermodynamic potential of said compound. Moreover, a  
direct correlation between the thermodynamic potential  
and the degree of saturation (DS) can often be assumed.  
15 Therefore, the equation

$$DS = \frac{\text{permeation rate}}{\text{permeation rate at saturation}}$$
  
was therefore assumed to be valid when estimating the DS  
values.

20 The  $t_p$ -values for the compositions Y1-Y4 according to  
the present invention were then investigated in the same  
manner as described above. These investigations showed  
that the  $t_p$  value for all the compositions Y1-Y4 exceeds  
6 weeks. At the time of filing the present application,  
25 no precipitation had yet been observed. Indeed, the  
precipitation preventing properties of the carrier matrix  
according to the present invention were clearly  
substantiated, particularly in comparison with the  $t_p$   
values depicted in Table 1 above.

Example 3; further evidence of increased thermodynamic potential attained in accordance with the present invention:

5 These experiments were performed in order to further substantiate the degree of saturation of metronidazole in the compositions X1-X4 and Y1-Y4. The Franz diffusion cell experiments were performed under the same conditions as in example 1 (*vide supra*), and the results are shown in Diagram 2.

10




Diagram 2. Amount of metronidazole permeated from compositions X1-X4 and Y1-Y4.

15 As depicted above, Diagram 2 shows that the chemical operation subjected to the compositions X1-X4 upon manufacturing of the compositions Y1-Y4 resulted in an increased thermodynamic potential of metronidazole, as is directly evidenced through the increased permeation rate.

20 The permeation rate for a Y composition has increased approximately 40% in comparison with its corresponding X composition.

In summary, it is clearly realised that biologically active compositions which are prepared or obtainable in accordance with the present invention are useful as medicaments. Furthermore, the biologically active compositions according to the invention are also useful in a non-medicinal context, such as in cosmetic skin products. More specifically, said compositions should be highly efficient in dermal application to a mammal, preferably man, as well as in any general application where a biological barrier is to be penetrated by a biologically active agent.

CLAIMS

1. A biologically active composition comprising a biologically active agent to be released therefrom, said biologically active agent being dissolved and/or  
5 dispersed in a carrier therefor, wherein said carrier is a liquid and/or solid non-crystalline matrix in which said biologically active agent is present in a supersaturated state, said supersaturated state being obtainable by subjecting one or more carrier starting  
10 substance(s) to such chemical operation(s) that said liquid and/or solid non-crystalline carrier matrix is provided in which the degree of saturation of said biologically active agent is higher than in said carrier starting substance(s), the biologically active agent  
15 being added before said chemical operation(s) has (have) been completed.

2. A composition according to claim 1, wherein said higher degree of saturation is the result of such  
20 chemical operation(s) that the solubility of the biologically active agent in said matrix is lower than the solubility thereof in said carrier starting substance(s).

25 3. A composition according to any one of claims 1 and 2, wherein said higher degree of saturation is the result of such chemical operation(s) that the degree of dissociation, aggregation and/or degree of protonation of the biologically active agent is different from the  
30 degree of dissociation, aggregation and/or degree of protonation of said agent in said carrier starting substance(s).

4. A composition according to any one of claims 1-3, wherein said biologically active agent is added before said chemical operation(s) has (have) been initiated.

5        5. A composition according to any one of claims 1-3, wherein said biologically active agent is added at a predetermined point of time after said chemical operation(s) has (have) been initiated, the composition thus obtained then being further subjected to said  
10 chemical operation(s).

6. A composition according to claim 5, wherein said predetermined point of time is from 1 minute to 6 months, preferably from 0,5 hours to 4 months after said chemical  
15 operation(s) has (have) been initiated.

7. A composition according to claim 6, wherein the composition is further subjected to said chemical operation(s) for a time period of about from 1 minute to  
20 6 months, preferably from 0,5 hours to 4 months.

8. A composition according to any one of claims 1-7, wherein said starting substance(s), or said formed non-crystalline matrix, act(s) as a solvent or dispersing  
25 medium.

9. A composition according to any one of claims 1-8, wherein said biologically active agent is added as a solid and/or liquid which is subsequently dissolved in  
30 said carrier.

10. A composition according to any one of claims 1-8, wherein said biologically active agent is added in the form of a solution or dispersion.

11. A composition according to any one of claims 1-10, wherein said biologically active agent is added above or around room temperature.

5        12. A composition according to any one of claims 1-11, wherein said chemical operation(s) comprise one or more chemical reactions.

10        13. A composition according to claim 12, wherein said chemical reaction(s) comprise etherifying, esterifying, hydrolysis, substitution, addition, elimination, oligomerising and/or polymerising reactions.

15        14. A composition according to claim 13, wherein said chemical reaction(s) is (are) selected and performed so as to provide optimal delivery rate of said biologically active agent.

20        15. A composition according to any one of claims 1-14, wherein said chemical operation(s) involve(s) subjecting said carrier starting substance(s) to a temperature of from around -50°C to around 300°C, preferably around 0-150°C.

25        16. A composition according to any one of claims 1-15, wherein said chemical operation(s) is (are) conducted for a time period of from 1 minute to 6 months, preferably from 0,5 hours to 4 months.

30        17. A composition according to any one of claims 1-16, wherein said carrier starting substance, or mixture of two or more different carrier starting substances, is selected from monomers, acids, such as mono-, di- or triacids or higher acids, alcohols, including mono-, di-  
35        or triols, ketones, aldehydes, amines, amides, anhydrides, lactides, glycolides, saccharides and derivatives thereof, acrylic or acrylamide type

compounds, such as methyl methacrylate, monomers of PEO-diacrylate, cyanoacrylate, acrylate saccharides, including acrylate starch, acrylate lactate, acrylate glycolate, isocyanates, ethylene oxide, propylene oxide, pyrrolidone, PEO-diacrylate, ethylene-vinyl acetate, monomers of organic siloxanes, and oligomers, polymers or prepolymers thereof.

18. A composition according to claim 17, wherein the acid is a monomeric acid and the alcohol is a monomeric alcohol, said non-crystalline matrix comprising an ester and/or polyester thereof.

19. A composition according to claim 18, wherein said monomeric acid is citric acid.

20. A composition according to any one of claims 18 and 19, wherein said monomeric alcohol is propylene glycol.

20

21. A composition according to any one of the preceding claims, which consists of one liquid or solid phase only.

22. A composition according to any one of the preceding claims, wherein the biologically active agent is a pharmaceutically active agent.

23. A composition according to claim 22, wherein the pharmaceutically active agent is selected from the group consisting of guanosides, corticosteroids, psychopharmaceutical hormones, oxicams, peptides, proteins, antibiotics, antivirals, antimicrobials, anticancer agents, antifungals, oestrogens, antiinflammatory agents, neuroleptic agents, melanocyte stimulants and gland stimulants, preferably stimulators



of sebaceous and pilo-sebaceous glands, and agents with an effect on mast cell secretion.

24. A composition according to any one of claims 22  
5 and 23 for use as a medicament.


25. A composition according to any one of the  
preceeding claims for topical, preferably dermal  
application to a mammal, preferably man.

10

26. A method for the preparation of a biologically  
active composition comprising a biologically active agent  
dissolved and/or dispersed in a carrier therefor, wherein  
a carrier starting substance, or a mixture of two or  
15 more different carrier starting substances, is (are)  
subjected to such chemical operation(s) that a liquid  
and/or solid non-crystalline carrier matrix is formed, in  
which the degree of saturation of said biologically  
active agent is higher than in said carrier starting  
20 substance(s), said biologically active agent being added  
before said chemical operation(s) has (have) been  
completed and in an amount such that a supersaturated  
state is obtained.

25 27. A method according to claim 26, wherein said  
composition is as defined in any one of claims 2-25.

Diagram 1



## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/SE 99/00824

## A. CLASSIFICATION OF SUBJECT MATTER

IPC6: A61K 9/00, A61K 9/70  
According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 9700670 A1 (BIOGLAN AB), 9 January 1997 (09.01.97) --	1-27
A	WO 9710812 A2 (CYGNUS, INC.), 27 March 1997 (27.03.97) --	1-27
A	GB 2306885 A (RECKITT & COLMAN PRODUCTS LIMITED), 14 May 1997 (14.05.97) --	1-27
A	US 4940701 A (ADRIAN F. DAVIS), 10 July 1990 (10.07.90) -- -----	1-27



Further documents are listed in the continuation of Box C.



See patent family annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

6 Sept 1999

Name and mailing address of the ISA/

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Date of mailing of the international search report

07 -09- 1999

Authorized officer

Eva Johansson/EÖ

Telephone No. +46 8 782 25 00

# INTERNATIONAL SEARCH REPORT

Information on patent family members

02/08/99

International application No.

PCT/SE 99/00824

Patent document cited in search report			Publication date	Patent family member(s)		Publication date
WO	9700670	A1	09/01/97	AU	695622 B	20/08/98
				AU	6247396 A	22/01/97
				CA	2225286 A	09/01/97
				DE	833611 T	07/01/99
				EP	0833611 A	08/04/98
				ES	2118677 T	01/10/98
				NZ	311208 A	29/03/99
				SE	9502244 D	00/00/00
WO	9710812	A2	27/03/97	AU	6968296 A	09/04/97
				CA	2229184 A	27/03/97
				EP	0848608 A	24/06/98
				US	5906830 A	25/05/99
GB	2306885	A	14/05/97	AU	7315596 A	29/05/97
				EP	0862417 A	09/09/98
				GB	9522885 D	00/00/00
				WO	9717062 A	15/05/97

**INTERNATIONAL SEARCH REPORT**  
Information on patent family members

02/08/99

International application No.  
PCT/SE 99/00824

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4940701 A	10/07/90	AT 71519 T	15/02/92
		AU 576344 B	25/08/88
		AU 3800885 A	01/08/85
		CA 1249520 A	31/01/89
		DE 3585162 A	27/02/92
		DK 30585 A	26/07/85
		DK 163799 B,C	06/04/92
		EP 0151953 A,B	21/08/85
		SE 0151953 T3	
		IE 57733 B	07/04/93
		JP 1741096 C	15/03/93
		JP 4030923 B	25/05/92
		JP 60174715 A	09/09/85
		US 4767751 A	30/08/88
		AT 96021 T	15/11/93
		AU 609756 B	09/05/91
		AU 8227587 A	16/06/88
		CA 1302256 A	02/06/92
		DE 3787881 D,T	10/02/94
		DK 649487 A	12/06/88
		EP 0272045 A,B	22/06/88
		SE 0272045 T3	
		HK 1006280 A	00/00/00
		IE 60778 B	10/08/94
		JP 2794285 B	03/09/98
		JP 63179821 A	23/07/88

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**